

sider it the drug of choice, now to be used in preference to paraldehyde, especially in children. For severe and intractable seizures, intravenous slow drip of 100 mg and 500 ml of normal saline solution is recommended with the total dose initially being 10 to 15 mg per square meter of body surface. The most significant side effects are moderate hypotension and mild respiratory depression, both most commonly seen in adults and in conjunction with barbiturates or paraldehyde. Diazepam has also proved useful as an oral adjunct with other drugs in the long-term control of convulsive disorders.

TOM W. ROBINSON, M.D.

REFERENCES

- Parsonage MJ: Use of diazepam in the treatment of severe convulsive status epilepticus. *Brit Med J* 5:85-88, 1967
 Calderon-Gonzales R, Mireles-Gonzales A: Management of prolonged motor seizure activity in children. *JAMA* 204:544-546, 1968
 Sawyer GT: Treatment of uncontrolled seizure activity with diazepam. *JAMA* 203:913-918, 1968
 Bailey DW, Fenichel GM: The treatment of prolonged seizure activity with intravenous diazepam. *J Pediatr* 73:923-927, 1968
 Bowe J: Status epilepticus and diazepam. *Brit Med J* 1:439-440, 1969

Growth Retardation in Environmental Deprivation

Environmental deprivation is perhaps the most frequent cause of growth failure in infancy. In older children it is less frequent as a cause of growth failure. If careful history taking, careful physical examination and simple laboratory testing (blood cell count, urinalysis, serum urea, protein-bound iodine) fail to reveal an organic cause, environmental deprivation should be considered. Infants with mental retardation may not grow adequately whether they are deprived or not. Admission to a hospital may be necessary to determine if an infant can gain weight if adequately nourished and cared for. Testing should be held to a minimum during this time.

S. A. KAPLAN, M.D.

REFERENCES

- Riley RL, Landwirth J, Kaplan SA, et al: Failure to thrive: An analysis of 83 cases. *Calif Med* 108:32-38, 1968
 Patton RG, Gardner LI: *Growth Failure in Maternal Deprivation*. Springfield, Illinois, Charles C Thomas, 1964

Intravenous Alimentation

Nearly complete alimentation can now be accomplished intravenously for long periods of time. This technique requires the placement of a silastic or polyvinyl catheter into the right atrium via the jugular, subclavian or femoral veins. In the right atrium, the large volume of blood flow will immediately dilute the hyperosmolar solution of 25 percent glucose, 4 percent fibrin hydrolysate, required electrolytes and vitamins. The volume of infusion is equivalent to the maintenance water requirements and should be given at a constant rate with a pump. The maximum rate of glucose utilization is 1.3 grams per kilogram of body weight per hour. A Millipore filter in the line will remove particulate matter and microorganisms. Essential fatty acids and trace metals are supplied by transfusion of plasma twice weekly. Improved survival has occurred in patients who are unable to eat because of prolonged ileus, chronic diarrhea, intestinal fistulae, peritonitis and sepsis. Complications include septicemia (particularly *Candida*), venous thrombosis and, rarely, pulmonary embolism. This technique of parenteral alimentation is one of the most important medical advances in many years.

ALFRED A. DELORIMIER, M.D.

REFERENCES

- Wilmore DW, Duderick SJ: Growth and development of an infant receiving all nutrients exclusively by vein. *JAMA* 203:860-864, 1968
 Filler RM, Eraklis AJ, Rubin VG, et al: Long-term parenteral nutrition in infants. *New Eng J Med* 281:589-594, 1969
 Duderick SJ, Wilmore DW, Vars HM, et al: Can intravenous feeding as the sole means of nutrition support growth of the child and restore weight loss in an adult? An affirmative answer. *Ann Surg* 169: 974-983, 1969

Drug Abuse

The most common mind-altering substances being abused by youth today are the hallucinogens, amphetamines and barbiturates. The hallucinogenic substances hemp (or marijuana), lysergic acid diethylamide (LSD), peyote, and mescaline all produce acute panic reactions, recurrent hallucinations, and serious psychotic change with varying frequency, but most commonly with LSD. Such untoward reactions with marijuana use are more rare due to the low content of the active

ingredient, tetrahydrocannabinol, in the material most commonly available. Methamphetamine, because of the rapidity with which dependence and tolerance develops and because abuse produces a chronic brain syndrome, is certainly the most dangerous drug being abused by young people. The oral use of dextroamphetamine, which is more common in high schools, is often followed by a shift to methamphetamine. Barbiturate use is also

common in the high school ages and represents an introductory move toward further experimentation with chemical solutions for life's problems.

HENRY B. BRUYN, M.D.

REFERENCES

- Drug Abuse Education—A Guide for the Professions. American Pharmaceutical Corporation, 1969
 Drug Dependence: A Guide for Physicians. Chicago, American Medical Association, 1969
 Blum RH, and associates: Students and Drugs. San Francisco, Jossey-Bass, 1969

ADVISORY PANEL SECTION ON PEDIATRICS

EUGENE GETTELMAN, M.D., Chairman

Encino

Chairman of CMA Section on Pediatrics

WILLIAM B. BELFORD, M.D.

*CMA Scientific Board
San Diego*

ROBERT F. CHINNOCK, M.D.

Loma Linda University School of Medicine

GEORGE DONNELL, M.D.

*CMA Scientific Board
Los Angeles*

MELVIN M. GRUMBACH, M.D.

University of California, San Francisco Medical Center

WILLIAM M. JENKINS, M.D.

*CMA Section on Pediatrics, Assistant Secretary
Oakland*

RUSSEL W. MAPES, M.D.

*American Academy of Pediatrics, President
Los Angeles*

ARTHUR J. MOSS, M.D.

*University of California, Los Angeles,
School of Medicine
Los Angeles*

THOMAS L. NELSON, M.D.

*University of California, Irvine,
California College of Medicine*

WILLIAM L. NYHAN, Ph.D.

*University of California, San Diego,
School of Medicine*

SAUL J. ROBINSON, M.D.

*American Academy of Pediatrics,
Chairman, District 9
San Francisco*

HERBERT SCHWARTZ, M.D.

Stanford University School of Medicine

ROBERT STEMPEL, M.D.

*University of California, Davis,
School of Medicine*

SHERROD C. SWIFT, M.D.

*CMA Scientific Board
Whittier*

CHESTER TANCREDI, M.D.

*CMA Section on Pediatrics, Secretary
San Diego*

PAUL F. WEHRLE, M.D.

*University of Southern California,
School of Medicine
Los Angeles*

In addition, the following physicians met with the panel and participated in the selection and preparation of these epitomes: GLENN E. AUSTIN, M.D., *American Academy of Pediatrics, Council on Pediatric Practice, Los Altos*; LEO S. BELL, M.D., *American Academy of Pediatrics, Council on Pediatric Practice, San Mateo*; JOHN A. BISHOP, M.D., *American Academy of Pediatrics, Chairman, Chapter 3, San Diego*; WILLIAM McCLELLAND BROWNLEE, M.D., *American Academy of Pediatrics, Treasurer, Chapter 3, San Diego*; RALPH W. COFFELT, M.D., *American Academy of Pediatrics, Immediate Past President, District 9, Burbank*; CARL W. ERICSON, M.D., *American Academy of Pediatrics, Pasadena*; BURTON FINK, M.D., *President, Los Angeles County Pediatric Society*; BIRT HARVEY, M.D., *American Academy of Pediatrics, Alternate Chairman, Chapter 1, Palo Alto*; ROBERT J. HARVEY, M.D., *CMA Section on Pediatrics, Immediate Past Chairman, San Francisco*; ALEXANDER HATOFF, M.D., *American Academy of Pediatrics, Chairman, Chapter 1, Oakland*; H. JAMES HOLROYD, M.D., *American Academy of Pediatrics, Secretary-Treasurer, Chapter 2, La Canada*; IAN BRUCE JOHNSTON, M.D., *American Academy of Pediatrics, Immediate Past Chairman, Chapter 1, Walnut Creek*; FREMONT P. KOCH, M.D., *American Academy of Pediatrics, Chairman, Chapter 2, Arcadia*; VICTOR Y. LINDBLADE, M.D., *American Academy of Pediatrics, Alternate Chairman, Chapter 3, San Diego*; WILLIAM D. MISBACH, M.D., *American Academy of Pediatrics, Alternate Chairman, Chapter 2, Encino*; EDWARD J. OTTENHEIMER, M.D., *American Academy of Pediatrics, Secretary, Chapter 3, San Diego*; TOM W. ROBINSON, M.D., *CMA Committee on School and College Health, Chairman, Newport Beach*; MELVIN H. SCHWARTZ, M.D., *American Academy of Pediatrics, Secretary, Chapter 1, Alameda*; MAXINE SEHRING, M.D., *School Health Committee, American Academy of Pediatrics, Walnut Creek*; EDWARD B. SHAW, M.D., *University of California, San Francisco Medical Center*; LOUISE YEAZELL, M.D., *American Academy of Pediatrics, Treasurer, Chapter 1, San Francisco*.